

Upali Amarasinghe ORCID iD: 0000-0002-0465-6950

Comparison of trophic structure of fish assemblages in two tropical streams in Sri Lanka: a seasonal dry zone stream and a perennial wet zone stream

Wasantha S. Weliange¹ | Upali S. Amarasinghe^{2*} | Jacobus Vijverberg³ | Maria Leichtfried⁴ | Leopold Füreder⁵

¹Department of Biological Sciences, Faculty of Applied Sciences, Rajarata University of Sri Lanka, Mihintale 50300, Sri Lanka.

²Department of Zoology and Environmental Management, University of Kelaniya, Kelaniya 11600, Sri Lanka

³Department of Aquatic Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Droevendaalsesteeg 10, 6708 PB Wageningen, The Netherlands

⁴Institute for Limnology of the Austrian Academy of Sciences. Mondseestr. 9, A-5310 Mondsee, Austria

⁵River Ecology and Conservation, Institute of Ecology, University of Innsbruck, Technikerstr. 25, A-6020 Innsbruck, Austria

* **Correspondence:** Prof. Upali S. Amarasinghe, Department of Zoology and Environmental Management, University of Kelaniya, Kelaniya 11600, Sri Lanka

E-mail: zoousa@kln.ac.lk; Fax: +94(0)-112-9144-79

Running title: Trophic structure of fish in two tropical streams

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/iroh.201601867.

This article is protected by copyright. All rights reserved.